

Contractile Units in Disordered Actomyosin Bundles Arise from F-Actin Buckling

Martin Lenz,¹ Todd Thoresen,² Margaret L. Gardel,^{1,2,3} and Aaron R. Dinner^{1,2,4,*}

¹James Franck Institute, University of Chicago, Chicago, Illinois 60637, USA

²Institute for Biophysical Dynamics, University of Chicago, Chicago, Illinois 60637, USA

³Department of Physics, University of Chicago, Chicago, Illinois 60637, USA

⁴Department of Chemistry, University of Chicago, Chicago, Illinois 60637, USA

(Received 19 October 2011; published 8 June 2012)

Bundles of filaments and motors are central to contractility in cells. The classic example is striated muscle, where actomyosin contractility is mediated by highly organized sarcomeres which act as fundamental contractile units. However, many contractile bundles *in vivo* and *in vitro* lack sarcomeric organization. Here we propose a model for how contractility can arise in bundles without sarcomeric organization and validate its predictions with experiments on a reconstituted system. In the model, internal stresses in frustrated arrangements of motors with diverse velocities cause filaments to buckle, leading to overall shortening. We describe the onset of buckling in the presence of stochastic motor head detachment and predict that buckling-induced contraction occurs in an intermediate range of motor densities. We then calculate the size of the “contractile units” associated with this process. Consistent with these results, our reconstituted actomyosin bundles show contraction at relatively high motor density, and we observe buckling at the predicted length scale.

DOI: 10.1103/PhysRevLett.108.238107

PACS numbers: 87.16.Ln, 87.16.A–, 87.16.Ka, 87.16.Nn

Contractility arising from interactions between myosin molecular motors and polar actin filaments (F-actin) is used ubiquitously by cells to build tension and drive morphological changes [1]. Such force transmission from molecular to cellular length scales is well understood in striated muscle, where it critically relies on highly organized structures known as sarcomeres [2]. In sarcomeres, myosin motors are restricted to the pointed end of F-actin, while passive actin cross-linkers are present at the barbed end. This arrangement is crucial to their contraction mechanism, as shown in Fig. 1(a). However, many contractile actomyosin bundles found *in vivo*, such as smooth muscle fibers [3], graded polarity bundles [4], and the contractile ring [5], lack a sarcomeric organization. Most recently, we have shown that *in vitro* bundles lacking apparent sarcomeric organization can also contract [6] [e.g., Fig. 1(b)]. In these disparate systems, contraction occurs with a well-defined contraction velocity per unit length, suggesting that contractile bundles can be meaningfully divided into elementary units that are arranged in series [5–7]. The mechanisms giving rise to such units in the absence of sarcomeric organization are not understood.

Much theoretical work on nonsarcomeric actomyosin assemblies posits contractility as a fundamental assumption, and predicts larger-scale effects such as polarity organization [9], the appearance of topological defects [10], active stiffening [11], and oscillatory behavior in cells [12]. Models that address the microscopic origin of contractility assume that myosin motors dwell at the barbed ends of F-actin, thus acting as transient static cross-linkers [13]. This generates sufficient sarcomerelike organization to elicit contraction [14]. Experimental evidence for this

behavior is unfortunately lacking, and it is thus important to investigate alternative routes to contractility.

In considering such mechanisms, it is important to recognize that actomyosin interactions can *a priori* elicit extension just as well as contraction. Figure 1(c) illustrates this using two elementary bundles, each made of two polar filaments—representing F-actin—and one motor—representing a whole myosin thick filament, itself comprising numerous individual myosin heads; we use this definition of a “motor” throughout. These elementary bundles contract when the motor is located in the vicinity of the filament pointed ends, but extend when it is close to the barbed ends. Overall contractility in nonsarcomeric bundles requires that the symmetry between these two competing tendencies be broken. We show in Ref. [15] that this necessitates (1) a dispersion of unloaded velocities to be present among the motors (as observed experimentally [16]) and (2) an asymmetric response of the filaments to longitudinally applied stresses, e.g., a tendency to yield under compression while resisting extension.

In this Letter, we use theory and experiments to demonstrate a mechanism for nonsarcomeric contractility compatible with these constraints. We first show experimentally that contraction in reconstituted actomyosin bundles is accompanied by F-actin buckling, an instance of the asymmetric filament response discussed above. We then investigate the general consequences of asymmetric filament response theoretically by considering the buildup of forces in a bundle with randomly arranged motors. We predict that buckling yields contraction, and occurs in an intermediate range of motor density. We also calculate a characteristic length scale between two buckles, which




FIG. 1. Contraction in actomyosin bundles. (a) Sarcomeric structure as in striated muscle. As motors tend to move toward the filament barbed ends, the sarcomeric structure imposes that each contractile unit (sarcomere) contracts. (b) Bundle devoid of sarcomeric organization or passive cross-linkers, as in our experiments. (c) Motors and polar filaments induce local contraction or extension depending on the geometry of their assembly (filament polarity always dictates the direction of motion [8]). (d) Time-lapse images of a bundle comprised of F-actin and fluorescent myosin thick filaments (inverted contrast) with $\ell_0 = 540$ nm. The initially wavy bundle becomes taut following the addition of 1 mM ATP at $t = 0$ s, indicating contraction. Scale bar, $5 \mu\text{m}$. (e) Similar experiment with $\ell_0 = 1.5 \mu\text{m}$, showing no contraction. Scale bar as in (d). See also movie S1 [18]. (f) Bundle contraction as a function of ℓ_0 . Bars indicate standard deviation ($n \geq 25$).

provides a natural size for a contractile unit. These predictions are consistent with experimental observations, suggesting that buckling underlies contractility in nonsarcomeric actomyosin bundles.

To form reconstituted actomyosin bundles, we follow the protocol described in Ref. [6]. We incubate F-actin with length $\ell_f \approx 5 \mu\text{m}$ with smooth muscle myosin thick filaments of length ≈ 300 nm in a buffer lacking adenosine triphosphate (ATP) such that thick filaments cross-link F-actin with high affinity. In this system, the dispersion of motor velocities necessary for contraction likely arises from the variation in number of myosin heads in the thick filaments. While flexible motors have been considered as a basis for contraction [17], this is unlikely to apply here as thick filaments are significantly more rigid than F-actin. The bundle lengths range from 10 to $100 \mu\text{m}$ with 4–6 F-actin per bundle cross-section, and no sarcomeric organization is observed. By varying the concentration of myosin filaments, the average spacing ℓ_0 between two consecutive myosin filaments can be varied from 390 nm to $5.3 \mu\text{m}$.

Once the bundles are formed, we perfuse a buffer containing 1 mM ATP, which causes bundles formed with high

myosin density ($\ell_0 = 540$ nm) to shorten by $\approx 10\%$ rapidly ($100\text{--}600 \text{ nm} \cdot \text{s}^{-1}$) [Fig. 1(d) and movie S1 [18]]. In contrast, contraction does not occur at low myosin density ($\ell_0 = 1.5 \mu\text{m}$) [Fig. 1(e) and movie S1 [18]]. A sharp transition between those two behaviors is observed at $\ell_0 = 1.3 \mu\text{m}$ [Fig. 1(f)].

To better understand the underpinnings of contractility in this system, we next examine individual bundles and observe F-actin buckling coincident with contraction [Fig. 2(a); movie S2 [18]]. Prior to ATP addition, compact bundles with aligned F-actin are observed. Upon ATP addition, the frequency of buckles increases rapidly during contraction, and then diminishes once contraction stops [Fig. 2(b)]. These F-actin buckles are dynamic, with their amplitude, curvature, and location changing over time.

Qualitatively, the relationship between buckling and contraction can be understood as follows. Consider two antiparallel filaments interacting through several different motors with distinct speeds [Fig. 2(c)]. As motors start to move relative to the filaments, stresses build in sections of the filament flanked by motors with different speeds. When the flanking motor proximal to the barbed end is faster than that proximal to the pointed end, compression arises. When it is slower, tension arises. Filament buckling breaks the symmetry between these respective tendencies to contraction and extension. Indeed, following buckling of the compressed filament sections, fast motors are free to move quickly while the others move slowly. This results in the growth of the compressed sections and shrinkage of the extended ones, and thus in overall bundle contraction




FIG. 2 (color online). Buckling in nonsarcomeric contractile actomyosin bundles. (a) Time-lapse images of fluorescent actin (inverted contrast) showing F-actin buckling (arrowheads) following the addition of 1 mM ATP at $t = 0$ s. Scale bar, $5 \mu\text{m}$. See also movie S2 [18]. (b) Relative contraction (filled squares) and number of F-actin buckles (open circles) as a function of time. Data show mean \pm sd averaged over $n = 3$ bundles with $\ell_0 \approx 1 \mu\text{m}$. (c) The presence of fast (gray) and slow (white) motors generically induce compressive (solid red) and extensile (hatched blue) stresses in filaments. (d) Buckling of the compressed filaments leads to an overall shortening of the bundle.

[Fig. 2(d)]. The region centered around each buckle thus plays the role of a contractile unit, whose typical size is equal to the distance ℓ_B between two buckles.

In this picture, the contractile behavior of the bundle hinges on the ability of the motors to induce filament buckling. At high motor density, we expect the bundle to be so strongly cross-linked that buckling becomes impossible despite the sizable stresses induced by a large number of motors. At low motor density, we expect that stochastic detachment of the motors undermines stress buildup and thus prevents buckling. Here we present a mathematical model to predict the range of myosin densities enabling buckling and the contractile unit length ℓ_B . These results are then compared with the observations in Figs. 1 and 2 to validate the proposed contraction mechanism.

The key assumptions of our model are that (1) motors have a dispersion in their unloaded velocities, (2) a section of filament between two motors buckles above a certain threshold force F_B , and (3) motors intermittently detach from the filaments, thus allowing local stress relaxation. We consider a bundle of weakly deformed filaments and ask whether the forces developing within it are sufficient to induce buckling [Fig. 3(a)].

To this end, we focus on a single filament of length ℓ_f and approximate its surroundings by an effective medium composed of evenly spaced pointlike motors separated by a distance $\ell_0 \ll \ell_f$ [Fig. 3(b)]. This divides the filament into discrete sections, which we label by $i = 0, \dots, \ell_f/\ell_0$. We take into account the possibility that the filaments are not straight, but bend away from the x axis, implying that the contour length L_i of filament section i can be larger than ℓ_0 . Defining f_i as the tension of filament section i ($f_i < 0$ for a compressed filament section), we expand its force-extension relationship for small deformations,

$$L_i = L_i(f_i = 0) - cf_i, \quad (1)$$




FIG. 3 (color online). Stress buildup in bundles with nonidentical motors. (a) In a bundle with motors having nonidentical velocities (shades of gray), filaments of lengths $\approx \ell_f$ are subjected to random motor forces at points $\approx \ell_0$ apart distributed throughout their length. (b) Prior to contraction, the environment of a filament of interest (red, delimited by dotted lines) can be approximated by a collection of evenly spaced motors (shades of gray).

where $c > 0$ is the filament compliance. We refer to the motor flanked by filament sections $i - 1$ and i as “motor i ,” and describe its operation by the simplified force-velocity relationship

$$f_{i-1} - f_i = F_i - \chi v_i. \quad (2)$$

Here v_i denotes the local velocity of the filament at the location of motor i and $\chi > 0$ is the motor susceptibility. Equations (1) and (2) yield a local relaxation time scale $\tau_r = \chi c/2$. The time-independent stall force of motor i is denoted by F_i in Eq. (2), and is drawn from a random distribution satisfying

$$\bar{F}_i = F_S \quad \text{and} \quad \overline{F_i F_j} - \bar{F}_i \bar{F}_j = \delta F_S^2 \delta_{ij}, \quad (3)$$

where bars denote averages over the motor distribution.

As a result, different motors have different unloaded velocities F_i/χ as required for contraction. Owing to the conservation of filament mass,

$$\frac{dL_i}{dt} = v_i - v_{i+1}. \quad (4)$$

Finally, a motor bound to several filaments as in Fig. 3(a) can transiently detach from one while still holding onto the others. We thus let each motor i randomly detach from the filament with a constant rate $1/\tau_d$. Following detachment, local filament stresses relax instantaneously, yielding $f_i = f_{i-1} = (f_i + f_{i-1})/2$. The motor then reattaches after a time much shorter than τ_r and τ_d . Since a motor in a dense bundle is typically close to several filaments, the probability that it detaches from all filaments at the same time and leaves the bundle is negligible. We denote by $\langle \dots \rangle$ the average over the Poisson process of motor detachment.

We obtain the space and time evolution of the filament tension $f(x, t)$ in the continuum limit $i \rightarrow x/\ell_0$ by combining Eqs. (1)–(4) and averaging over motor detachment,

$$\partial_t \langle f \rangle - D \partial_x^2 \langle f \rangle = (\ell_0/2\tau_r) \partial_x F, \quad (5)$$

where $D = \ell_0^2(\tau_r^{-1} + \tau_d^{-1})/2$. The right-hand side of Eq. (5) involves the spatial gradient of the stall force $F(x)$, reflecting the fact that nonidentical motors lead to force buildup. This effect competes with the relaxation of filament forces through motor detachment, which enters through the diffusion term $D \partial_x^2 \langle f \rangle$.

An initially relaxed filament [$f(x, t = 0) = 0$] experiences a vanishing average force $\langle \bar{f} \rangle(x, t) = 0$ throughout its dynamics. To quantify the magnitude of the motor-induced stress, we use Eqs. (1)–(4) and average over motor detachment to calculate the rms filament force in the continuum limit,

$$\left(\langle \bar{f}^2 \rangle \right)^{1/2} = f_\infty \left\{ \sum_{n \in \mathbb{Z}^*} \left[\frac{3}{n^2 \pi^2} (1 - e^{-n^2 D t / (2\pi \ell_f)^2})^2 \right] \right\}^{1/2}. \quad (6)$$

This force increases monotonically from zero at $t = 0$ to $f_\infty = (\ell_f/12\ell_0)^{1/2} \times \delta F_S / (1 + \tau_r/\tau_d)$ at $t = \infty$ [Figs. 4(a) and 4(b)]. We next estimate the dependence of the ratio

τ_r/τ_d on the experimentally accessible parameter ℓ_0 . A wormlike chain model for filament elasticity yields $c \approx \ell_0^4/k_B T \ell_p^2$, where ℓ_p is the filament persistence length [19], and we approximate $\chi \approx F_S/v$, where v is a characteristic motor velocity. This implies $\tau_r/\tau_d \approx (\ell_0/\ell_0^*)^{407.322510.4535712.9699m0c94.411c7.355417.326.72n0}$

We thank Yitzhak Rabin and Tom Witten for useful discussions. This work was supported by NSF DMR-MRSEC 0820054, NIH P50 GM081892, and NIH DP10D00354.

*dinner@uchicago.edu

- [1] J. Stricker, T. Falzone, and M. L. Gardel, *J. Biomech.* **43**, 9 (2010).
- [2] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, and P. Walter, *Essential Cell Biology* (Garland, New York, 1998).
- [3] F. S. Fay, K. Fujiwara, D. D. Rees, and K. E. Fogarty, *J. Cell Biol.* **96**, 783 (1983).
- [4] L. P. Cramer, M. Siebert, and T. J. Mitchison, *J. Cell Biol.* **136**, 1287 (1997).
- [5] Ana Carvalho, Arshad Desai, and Karen Oegema, *Cell* **137**, 926 (2009).
- [6] T. Thoresen, M. Lenz, and M. L. Gardel, *Biophys. J.* **100**, 2698 (2011).
- [7] W. M. Bement and D. G. Capco, *Cell Motil. Cytoskeleton* **20**, 145 (1991); A. M. Herrera, B. E. McParland, A. Bienkowska, R. Tait, P. D. Paré, and C. Y. Seow, *J. Cell Sci.* **118**, 2381 (2005).
- [8] J. R. Sellers and B. Kachar, *Science* **249**, 406 (1990).
- [9] K. Kruse, A. Zumdieck, and F. Jülicher, *Europhys. Lett.* **64**, 716 (2003); N. Yoshinaga, J.-F. Joanny, J. Prost, and P. Marcq, *Phys. Rev. Lett.* **105**, 238103 (2010).
- [10] K. Kruse, J. F. Joanny, F. Jülicher, J. Prost, and K. Sekimoto, *Phys. Rev. Lett.* **92**, 078101 (2004); *Eur. Phys. J. E* **16**, 5 (2005).
- [11] F. C. MacKintosh and A. J. Levine, *Phys. Rev. Lett.* **100**, 018104 (2008).
- [12] G. Salbreux, J. F. Joanny, J. Prost, and P. Pullarkat, *Phys. Biol.* **4**, 268 (2007).
- [13] K. Kruse and F. Jülicher, *Phys. Rev. Lett.* **85**, 1778 (2000); K. Kruse and K. Sekimoto, *Phys. Rev. E* **66**, 031904 (2002); K. Kruse and F. Jülicher, *Phys. Rev. E* **67**, 051913 (2003); T. B. Liverpool and M. C. Marchetti, *Phys. Rev. Lett.* **90**, 138102 (2003); R. Peter, V. Schaller, F. Ziebert, and W. Zimmermann, *New J. Phys.* **10**, 035002 (2008).
- [14] A. Zemel and A. Mogilner, *Phys. Chem. Chem. Phys.* **11**, 4821 (2009).
- [15] M. Lenz, M. L. Gardel, and A. R. Dinner, *New J. Phys.* **14**, 033037 (2012).
- [16] A. Yamada, N. Ishii, and K. Takahashi, *J. Biochem. (Tokyo)* **108**, 341 (1990).
- [17] T. B. Liverpool, M. C. Marchetti, J.-F. Joanny, and J. Prost, *Europhys. Lett.* **85**, 18007 (2009).
- [18] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.108.238107> for a study of nonidentical motor spacings and force-dependent filament detachment.
- [19] Theo Odijk, *Macromolecules* **28**, 7016 (1995).
- [20] B. Gilboa, D. Gillo, O. Farago, and A. Bernheim-Groswasser, *Soft Matter* **5**, 2223 (2009).
- [21] J. Berro, A. Michelot, L. Blanchoin, D. R. Kovar, and J.-L. Martiel, *Biophys. J.* **92**, 2546 (2007).
- [22] Y. Tanaka, A. Ishijima, and S. Ishiwata, *Biochim. Biophys. Acta* **1159**, 94 (1992).
- [23] M. Soares e Silva, M. Depken, B. Stuhmann, M. Korsten, F. C. MacKintosh, and G. H. Koenderink, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 9408 (2011).